


AMANTADINE- amantadine solution
Xttrium Laboratories, Inc

Amantadine 16oz, 10mL Lid, Insert

Description

Amantadine hydrochloride, USP is designated generically as amantadine hydrochloride and chemically as 1-adamantanamine hydrochloride.



$C_{10}H_{17}N \cdot HCl$

M.W. 187.71

Amantadine hydrochloride is a stable white or nearly white crystalline powder, freely soluble in water and soluble in alcohol and in chloroform.

Amantadine hydrochloride has pharmacological actions as both an anti-Parkinson and an antiviral drug.

Amantadine Hydrochloride Oral Solution, USP contains 50 mg of amantadine hydrochloride per 5 mL and has the following inactive ingredients: anhydrous citric acid, artificial raspberry flavor, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, sodium citrate dihydrate, and sorbitol solution.

Clinical Pharmacology

Pharmacodynamics

Mechanism of Action: Antiviral: The mechanism by which amantadine hydrochloride exerts its antiviral activity is not clearly understood. It appears to mainly prevent the release of infectious viral nucleic acid into the host cell by interfering with the function of the transmembrane domain of the viral M2 protein. In certain cases, amantadine hydrochloride is also known to prevent virus assembly during virus replication. It does not appear to interfere with the immunogenicity of inactivated influenza A virus vaccine.

Antiviral Activity: Amantadine hydrochloride inhibits the replication of influenza A virus isolates from each of the subtypes, i.e., H1N1, H2N2 and H3N2. It has very little or no activity against influenza B virus isolates. A quantitative relationship between the *in vitro* susceptibility of influenza A virus to amantadine hydrochloride and the clinical response to therapy has not been established in man. Sensitivity test results, expressed as the

concentration of amantadine hydrochloride required to inhibit by 50% the growth of virus (ED_{50}) in tissue culture vary greatly (from 0.1 $\mu\text{g/mL}$ to 25.0 $\mu\text{g/mL}$) depending upon the assay protocol used, size of virus inoculum, isolates of influenza A virus strains tested, and the cell type used. Host cells in tissue culture readily tolerated amantadine hydrochloride up to a concentration of 100 $\mu\text{g/mL}$.

Drug Resistance: Influenza A variants with reduced *in vitro* sensitivity to amantadine hydrochloride have been isolated from epidemic strains in areas where adamantane derivatives are being used. Influenza viruses with reduced *in vitro* sensitivity have been shown to be transmissible and to cause typical influenza illness. The quantitative relationship between the *in vitro* sensitivity of influenza A variants to amantadine hydrochloride and the clinical response to therapy has not been established.

Mechanism of Action: Parkinson's Disease: The mechanism of action of amantadine hydrochloride in the treatment of Parkinson's disease and drug-induced extrapyramidal reactions is not known. Data from earlier animal studies suggest that amantadine hydrochloride may have direct and indirect effects on dopamine neurons. More recent studies have demonstrated that amantadine hydrochloride is a weak, non-competitive NMDA receptor antagonist ($K_i = 10\mu\text{M}$). Although amantadine hydrochloride has not been shown to possess direct anticholinergic activity in animal studies, clinically, it exhibits anticholinergic-like side effects such as dry mouth, urinary retention, and constipation.

Pharmacokinetics

Amantadine hydrochloride is well absorbed orally. Maximum plasma concentrations are directly related to dose for doses up to 200 mg/day. Doses above 200 mg/day may result in a greater than proportional increase in maximum plasma concentrations. It is primarily excreted unchanged in the urine by glomerular filtration and tubular secretion. Eight metabolites of amantadine hydrochloride have been identified in human urine. One metabolite, an N-acetylated compound, was quantified in human urine and accounted for 5 to 15% of the administered dose. Plasma acetylamantadine accounted for up to 80% of the concurrent amantadine hydrochloride plasma concentration in 5 of 12 healthy volunteers following the ingestion of a 200 mg dose of amantadine hydrochloride. Acetylamantadine was not detected in the plasma of the remaining seven volunteers. The contribution of this metabolite to efficacy or toxicity is not known. There appears to be a relationship between plasma amantadine hydrochloride concentrations and toxicity. As concentration increases, toxicity seems to be more prevalent, however, absolute values of amantadine hydrochloride concentrations associated with adverse effects have not been fully defined.

Amantadine hydrochloride pharmacokinetics were determined in 24 normal adult male volunteers after the oral administration of a single amantadine hydrochloride 100 mg soft gel capsule. The mean $\pm$ SD maximum plasma concentration was 0.22 ± 0.03 $\mu\text{g/mL}$ (range: 0.18 to 0.32 $\mu\text{g/mL}$). The time to peak concentration was 3.3 ± 1.5 hours (range: 1.5 to 8.0 hours). The apparent oral clearance was 0.28 ± 0.11 L/hr/kg (range: 0.14 to 0.62 L/hr/kg). The half-life was 17 ± 4 hours (range: 10 to 25 hours). Across other studies, amantadine hydrochloride plasma half-life has averaged 16 ± 6 hours (range: 9 to 31 hours) in 19 healthy volunteers.

After oral administration of a single dose of 100 mg amantadine hydrochloride in a syrup formulation to five healthy volunteers, the mean $\pm$ SD maximum plasma concentration C_{max} was 0.24 ± 0.04 $\mu\text{g/mL}$ and ranged from 0.18 to 0.28 $\mu\text{g/mL}$. After 15 days of amantadine hydrochloride 100 mg b.i.d., the C_{max} was 0.47 ± 0.11 $\mu\text{g/mL}$ in four of the five volunteers. Across studies, the time to C_{max} (T_{max}) averaged about 2 to 4 hours. Plasma amantadine hydrochloride clearance ranged from 0.2 to 0.3 L/hr/kg after the

administration of 5 mg to 25 mg intravenous doses of amantadine hydrochloride to 15 healthy volunteers.

In six healthy volunteers, the ratio of amantadine hydrochloride renal clearance to apparent oral plasma clearance was 0.79 ± 0.17 (mean $\pm$ SD).

The volume of distribution determined after the intravenous administration of amantadine hydrochloride to 15 healthy subjects was 3 to 8 L/kg, suggesting tissue binding. Amantadine hydrochloride, after single oral 200 mg doses to 6 healthy young subjects and to 6 healthy elderly subjects has been found in nasal mucus at mean $\pm$ SD concentrations of 0.15 ± 0.16 , 0.28 ± 0.26 , and 0.39 ± 0.34 $\mu\text{g/g}$ at 1, 4, and 8 hours after dosing, respectively. These concentrations represented $31 \pm 33\%$, $59 \pm 61\%$, and $95 \pm 86\%$ of the corresponding plasma amantadine hydrochloride concentrations.

Amantadine hydrochloride is approximately 67% bound to plasma proteins over a concentration range of 0.1 to 2.0 $\mu\text{g/mL}$. Following the administration of amantadine hydrochloride 100 mg as a single dose, the mean $\pm$ SD red blood cell to plasma ratio ranged from 2.7 ± 0.5 in 6 healthy subjects to 1.4 ± 0.2 in 8 patients with renal insufficiency.

The apparent oral plasma clearance of amantadine hydrochloride is reduced and the plasma half-life and plasma concentrations are increased in healthy elderly individuals age 60 and older. After single dose administration of 25 to 75 mg to 7 healthy, elderly male volunteers, the apparent plasma clearance of amantadine hydrochloride was 0.10 ± 0.04 L/hr/kg (range 0.06 to 0.17 L/hr/kg) and the half-life was 29 ± 7 hours (range 20 to 41 hours). Whether these changes are due to decline in renal function or other age related factors is not known.

In a study of young healthy subjects (n=20), mean renal clearance of amantadine hydrochloride, normalized for body mass index, was 1.5 fold higher in males compared to females ($p < 0.032$).

Compared with otherwise healthy adult individuals, the clearance of amantadine hydrochloride is significantly reduced in adult patients with renal insufficiency. The elimination half-life increases two to three fold or greater when creatinine clearance is less than 40 mL/min/1.73 m² and averages eight days in patients on chronic maintenance hemodialysis. Amantadine hydrochloride is removed in negligible amounts by hemodialysis.

The pH of the urine has been reported to influence the excretion rate of amantadine hydrochloride. Since the excretion rate of amantadine hydrochloride increases rapidly when the urine is acidic, the administration of urine acidifying drugs may increase the elimination of the drug from the body.

Indications and Usage

Amantadine Hydrochloride Oral Solution, USP is indicated for the prophylaxis and treatment of signs and symptoms of infection caused by various strains of influenza A virus. Amantadine hydrochloride is also indicated in the treatment of parkinsonism and drug-induced extrapyramidal reactions.

Influenza A Prophylaxis: Amantadine hydrochloride is indicated for chemoprophylaxis against signs and symptoms of influenza A virus infection. Because amantadine hydrochloride does not completely prevent the host immune response to influenza A infection, individuals who take this drug may still develop immune responses to natural disease or vaccination and may be protected when later exposed to antigenically related viruses. Following vaccination during an influenza A outbreak, amantadine hydrochloride prophylaxis should be considered for the 2- to 4-week time period required to develop

an antibody response.

Influenza A Treatment: Amantadine hydrochloride is also indicated in the treatment of uncomplicated respiratory tract illness caused by influenza A virus strains especially when administered early in the course of illness. There are no well-controlled clinical studies demonstrating that treatment with amantadine hydrochloride will avoid the development of influenza A virus pneumonitis or other complications in high risk patients.

There is no clinical evidence indicating that amantadine hydrochloride is effective in the prophylaxis or treatment of viral respiratory tract illnesses other than those caused by influenza A virus strains.

The following points should be considered before initiating treatment or prophylaxis with amantadine hydrochloride:

- Amantadine hydrochloride is not a substitute for early vaccination on an annual basis as recommended by the Centers for Disease Control and Prevention Advisory Committee on Immunization Practices.
- Influenza viruses change over time. Emergence of resistance mutations could decrease drug effectiveness. Other factors (for example, changes in viral virulence) might also diminish clinical benefit of antiviral drugs. Prescribers should consider available information on influenza drug susceptibility patterns and treatment effects when deciding whether to use amantadine hydrochloride.

Parkinson's Disease/Syndrome: Amantadine hydrochloride is indicated in the treatment of idiopathic Parkinson's disease (Paralysis Agitans), postencephalitic parkinsonism, and symptomatic parkinsonism which may follow injury to the nervous system by carbon monoxide intoxication. It is indicated in those elderly patients believed to develop parkinsonism in association with cerebral arteriosclerosis. In the treatment of Parkinson's disease, amantadine hydrochloride is less effective than levodopa, (-)-3-(3,4-dihydroxyphenyl)-L-alanine, and its efficacy in comparison with the anticholinergic antiparkinson drugs has not yet been established.

Drug-Induced Extrapyrmidal Reactions:

Amantadine hydrochloride is indicated in the treatment of drug-induced extrapyramidal reactions. Although anticholinergic-type side effects have been noted with amantadine hydrochloride when used in patients with drug-induced extrapyramidal reactions, there is a lower incidence of these side effects than that observed with the anticholinergic antiparkinson drugs.

Contraindications

Amantadine hydrochloride is contraindicated in patients with known hypersensitivity to amantadine hydrochloride or to any of the other ingredients in Amantadine Hydrochloride Oral Solution, USP.

WARNINGS

Deaths: Deaths have been reported from overdose with amantadine hydrochloride. The lowest reported acute lethal dose was 1 gram. Acute toxicity may be attributable to the anticholinergic effects of amantadine hydrochloride. Drug overdose has resulted in cardiac, respiratory, renal or central nervous system toxicity. Cardiac dysfunction includes arrhythmia, tachycardia and hypertension (see **OVERDOSAGE**).

Deaths due to drug accumulation (overdose) have been reported in patients with renal impairment, who were prescribed higher than recommended doses of amantadine hydrochloride for their level of renal function (see **DOSAGE AND ADMINISTRATION; Dosage of Impaired Renal Function** and **OVERDOSAGE**).

Suicide Attempts: Suicide attempts, some of which have been fatal, have been reported in patients treated with amantadine hydrochloride, many of whom received short courses for influenza treatment or prophylaxis. The incidence of suicide attempts is not known and the pathophysiologic mechanism is not understood. Suicide attempts and suicidal ideation have been reported in patients with and without prior history of psychiatric illness. Amantadine hydrochloride can exacerbate mental problems in patients with a history of psychiatric disorders or substance abuse. Patients who attempt suicide may exhibit abnormal mental states which include disorientation, confusion, depression, personality changes, agitation, aggressive behavior, hallucinations, paranoia, other psychotic reactions, and somnolence or insomnia. Because of the possibility of serious adverse effects, caution should be observed when prescribing amantadine hydrochloride to patients being treated with drugs having CNS effects, or for whom the potential risks outweigh the benefit of treatment.

CNS Effects: Patients with a history of epilepsy or other “seizures” should be observed closely for possible increased seizure activity.

Patients receiving amantadine hydrochloride who note central nervous system effects or blurring of vision should be cautioned against driving or working in situations where alertness and adequate motor coordination are important.

Other: Patients with a history of congestive heart failure or peripheral edema should be followed closely as there are patients who developed congestive heart failure while receiving amantadine hydrochloride.

Patients with Parkinson’s disease improving on amantadine hydrochloride should resume normal activities gradually and cautiously, consistent with other medical considerations, such as the presence of osteoporosis or phlebothrombosis.

Because amantadine hydrochloride has anticholinergic effects and may cause mydriasis, it should not be given to patients with untreated angle closure glaucoma.

Corneal Edema: Corneal edema has been reported in patients taking amantadine. Symptoms include sudden onset of blurry vision, or progressive vision loss, with or without eye pain. Corneal involvement is usually bilateral. Onset can occur from a few weeks to several years after starting amantadine. Resolution of symptoms typically begins within weeks of amantadine cessation. However, corneal grafts have been required in some patients when the condition is not recognized. Permanent damage can occur if amantadine is continued. Ask patients if their vision has changed and obtain ophthalmologic examinations to rule out corneal edema should vision changes occur after initiation of therapy with amantadine. If corneal edema occurs, taper and discontinue amantadine [see **DOSAGE AND ADMINISTRATION**].

Precautions

Amantadine hydrochloride should not be discontinued abruptly in patients with Parkinson's disease since a few patients have experienced a parkinsonian crisis, i.e., a sudden marked clinical deterioration, when this medication was suddenly stopped. The dose of anticholinergic drugs or of amantadine hydrochloride should be reduced if atropine-like effects appear when these drugs are used concurrently. Abrupt discontinuation may also precipitate delirium, agitation, delusions, hallucinations,

paranoid reaction, stupor, anxiety, depression and slurred speech.

Neuroleptic Malignant Syndrome (NMS): Sporadic cases of possible Neuroleptic Malignant Syndrome (NMS) have been reported in association with dose reduction or withdrawal of amantadine hydrochloride therapy. Therefore, patients should be observed carefully when the dosage of amantadine hydrochloride is reduced abruptly or discontinued, especially if the patient is receiving neuroleptics.

NMS is an uncommon but life-threatening syndrome characterized by fever or hyperthermia; neurologic findings including muscle rigidity, involuntary movements, altered consciousness; mental status changes; other disturbances such as autonomic dysfunction, tachycardia, tachypnea, hyper- or hypotension; laboratory findings such as creatine phosphokinase elevation, leukocytosis, myoglobinuria, and increased serum myoglobin.

The early diagnosis of this condition is important for the appropriate management of these patients. Considering NMS as a possible diagnosis and ruling out other acute illnesses (e.g., pneumonia, systemic infection, etc.) is essential. This may be especially complex if the clinical presentation includes both serious medical illness and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever and primary central nervous system (CNS) pathology.

The management of NMS should include: 1) intensive symptomatic treatment and medical monitoring, and 2) treatment of any concomitant serious medical problems for which specific treatments are available. Dopamine agonists, such as bromocriptine, and muscle relaxants, such as dantrolene are often used in the treatment of NMS, however, their effectiveness has not been demonstrated in controlled studies.

Renal disease: Because amantadine hydrochloride is mainly excreted in the urine, it accumulates in the plasma and in the body when renal function declines. Thus, the dose of amantadine hydrochloride should be reduced in patients with renal impairment and in individuals who are 65 years of age or older (see **DOSAGE AND ADMINISTRATION; Dosage for Impaired Renal Function**).

Liver disease: Care should be exercised when administering amantadine hydrochloride to patients with liver disease. Rare instances of reversible elevation of liver enzymes have been reported in patients receiving amantadine hydrochloride, though a specific relationship between the drug and such changes has not been established.

Impulse Control/Compulsive Behaviors:

Postmarketing reports suggest that patients treated with anti-Parkinson medications can experience intense urges to gamble, increased sexual urges, intense urges to spend money uncontrollably, and other intense urges. Patients may be unable to control these urges while taking one or more of the medications that are generally used for the treatment of Parkinson's disease and that increase central dopaminergic tone, including Amantadine hydrochloride. In some cases, although not all, these urges were reported to have stopped when the dose was reduced or the medication was discontinued.

Because patients may not recognize these behaviors as abnormal it is important for prescribers to specifically ask patients or their caregivers about the development of new or increased gambling urges, sexual urges, uncontrolled spending or other urges while being treated with Amantadine hydrochloride. Physicians should consider dose reduction or stopping the medication if a patient develops such urges while taking amantadine hydrochloride.

Melanoma:

Epidemiological studies have shown that patients with Parkinson's disease have a higher risk (2- to approximately 6-fold higher) of developing melanoma than the general

population. Whether the increased risk observed was due to Parkinson's disease or other factors, such as drugs used to treat Parkinson's disease, is unclear.

For the reasons stated above, patients and providers are advised to monitor for melanomas frequently and on a regular basis when using amantadine hydrochloride for any indication. Ideally, periodic skin examinations should be performed by appropriately qualified individuals (e.g., dermatologists).

Other: The dose of amantadine hydrochloride may need careful adjustment in patients with congestive heart failure, peripheral edema, or orthostatic hypotension. Care should be exercised when administering amantadine hydrochloride to patients with a history of recurrent eczematoid rash, or to patients with psychosis or severe psychoneurosis not controlled by chemotherapeutic agents.

Serious bacterial infections may begin with influenza-like symptoms or may coexist with or occur as complications during the course of influenza. Amantadine hydrochloride has not been shown to prevent such complications.

Information for Patients: Patients should be advised of the following information:

Blurry vision, eye pain, loss of vision, and/or impaired mental acuity may occur.

Gradually increase physical activity as the symptoms of Parkinson's disease improve.

Avoid excessive alcohol usage, since it may increase the potential for CNS effects such as dizziness, confusion, lightheadedness and orthostatic hypotension.

Avoid getting up suddenly from a sitting or lying position. If dizziness or lightheadedness occurs, notify physician.

Notify physician if mood/mental changes, swelling of extremities, difficulty urinating and/or shortness of breath occur.

Do not take more medication than prescribed because of the risk of overdose. If there is no improvement in a few days, or if medication appears less effective after a few weeks, discuss with a physician.

Consult physician before discontinuing medication. Seek medical attention immediately if it is suspected that an overdose of medication has been taken.

Drug Interactions: Careful observation is required when amantadine hydrochloride is administered concurrently with central nervous system stimulants.

Agents with anticholinergic properties may potentiate the anticholinergic-like side effects of amantadine hydrochloride.

Coadministration of thioridazine has been reported to worsen the tremor in elderly patients with Parkinson's disease, however, it is not known if other phenothiazines produce a similar response.

Coadministration of triamterene/hydrochlorothiazide resulted in a higher plasma amantadine hydrochloride concentration in a 61-year-old man receiving amantadine hydrochloride 100 mg TID for Parkinson's disease.¹ It is not known which of the components of triamterene/hydrochlorothiazide contributed to the observation or if related drugs produce a similar response.

Coadministration of quinine or quinidine with amantadine hydrochloride was shown to reduce the renal clearance of amantadine hydrochloride by about 30%.

The concurrent use of amantadine hydrochloride with live attenuated influenza vaccine (LAIV) intranasal has not been evaluated. However, because of the potential for interference between these products, LAIV should not be administered within 2 weeks before or 48 hours after administration of amantadine hydrochloride, unless medically indicated. The concern about possible interference arises from the potential for antiviral drugs to inhibit replication of live vaccine virus. Trivalent inactivated influenza vaccine can be administered at any time relative to use of amantadine hydrochloride.

Carcinogenesis, Mutagenesis: Long-term *in vivo* animal studies designed to evaluate

the carcinogenic potential of amantadine hydrochloride have not been performed. In several *in vitro* assays for gene mutation, amantadine hydrochloride did not increase the number of spontaneously observed mutations in four strains of *Salmonella typhimurium* (Ames Test) or in a mammalian cell line (Chinese Hamster Ovary cells) when incubations were performed either with or without a liver metabolic activation extract. Further, there was no evidence of chromosome damage observed in an *in vitro* test using freshly derived and stimulated human peripheral blood lymphocytes (with and without metabolic activation) or in an *in vivo* mouse bone marrow micronucleus test (140 to 550 mg/kg; estimated human equivalent doses of 11.7 to 45.8 mg/kg based on body surface area conversion).

Impairment of Fertility: The effect of amantadine hydrochloride on fertility has not been adequately tested, that is, in a study conducted under Good Laboratory Practice (GLP) and according to current recommended methodology. In a three litter, non-GLP, reproduction study in rats, amantadine hydrochloride at a dose of 32 mg/kg/day (equal to the maximum recommended human dose on a mg/m² basis) administered to both males and females slightly impaired fertility. There were no effects on fertility at a dose level of 10 mg/kg/day (or 0.3 times the maximum recommended human dose on a mg/m² basis); intermediate doses were not tested.

Failed fertility has been reported during human in vitro fertilization (IVF) when the sperm donor ingested amantadine hydrochloride 2 weeks prior to, and during the IVF cycle.

Pregnancy:

Teratogenic Effects : The effect of amantadine hydrochloride on embryofetal and perinatal development has not been adequately tested, that is, in studies conducted under Good Laboratory Practice (GLP) and according to current recommended methodology. However, in two non-GLP studies in rats in which females were dosed from 5 days prior to mating to Day 6 of gestation or on Days 7 to 14 of gestation, amantadine hydrochloride produced increases in embryonic death at an oral dose of 100 mg/kg (or 3 times the maximum recommended human dose on a mg/m² basis). In the non-GLP rat study in which females were dosed on Days 7 to 14 of gestation, there was a marked increase in severe visceral and skeletal malformations at oral doses of 50 and 100 mg/kg (or 1.5 and 3 times, respectively, the maximum recommended human dose on a mg/m² basis). The no-effect dose for teratogenicity was 37 mg/kg (equal to the maximum recommended human dose on a mg/m² basis). The safety margins reported may not accurately reflect the risk considering the questionable quality of the study on which they are based. There are no adequate and well-controlled studies in pregnant women. Human data regarding teratogenicity after maternal use of amantadine hydrochloride is scarce. Tetralogy of Fallot and tibial hemimelia (normal karyotype) occurred in an infant exposed to amantadine hydrochloride during the first trimester of pregnancy (100 mg P.O. for 7 days during the 6th and 7th week of gestation). Cardiovascular maldevelopment (single ventricle with pulmonary atresia) was associated with maternal exposure to amantadine hydrochloride (100 mg/d) administered during the first 2 weeks of pregnancy. Amantadine hydrochloride should be used during pregnancy only if the potential benefit justifies the potential risk to the embryo or fetus.

Nursing Mothers: Amantadine hydrochloride is excreted in human milk. Use is not recommended in nursing mothers.

Pediatric Use: The safety and efficacy of amantadine hydrochloride in newborn infants and infants below the age of 1 year have not been established.

Usage in the Elderly: Because amantadine hydrochloride is primarily excreted in the urine, it accumulates in the plasma and in the body when renal function declines. Thus,

the dose of amantadine hydrochloride should be reduced in patients with renal impairment and in individuals who are 65 years of age or older. The dose of amantadine hydrochloride may need reduction in patients with congestive heart failure, peripheral edema, or orthostatic hypotension (see **DOSAGE AND ADMINISTRATION**).

Adverse Reactions

The adverse reactions reported most frequently at the recommended dose of amantadine hydrochloride (5 to 10%) are: nausea, dizziness (lightheadedness), and insomnia.

Less frequently (1 to 5%) reported adverse reactions are:

depression, anxiety and irritability, hallucinations, confusion, anorexia, dry mouth, constipation, ataxia, livedo reticularis, peripheral edema, orthostatic hypotension, headache, somnolence, nervousness, dream abnormality, agitation, dry nose, diarrhea and fatigue.

Infrequently (0.1 to 1%) occurring adverse reactions are: congestive heart failure, psychosis, urinary retention, dyspnea, skin rash, vomiting, weakness, slurred speech, euphoria, thinking abnormality, amnesia, hyperkinesia, hypertension, decreased libido, and visual disturbance, including punctate subepithelial or other corneal opacity, corneal edema, decreased visual acuity, sensitivity to light, and optic nerve palsy.

Rare (less than 0.1%) occurring adverse reactions are:

instances of convulsion, leukopenia, neutropenia, eczematoid dermatitis, oculogyric episodes, suicidal attempt, suicide, and suicidal ideation (see **WARNINGS**).

Other adverse reactions reported during postmarketing experience with amantadine hydrochloride usage include:

Nervous System/Psychiatric: coma, stupor, delirium, hypokinesia, hypertonia, delusions, aggressive behavior, paranoid reaction, manic reaction, involuntary muscle contractions, gait abnormalities, paresthesia, EEG changes, and tremor. Abrupt discontinuation may also precipitate delirium, agitation, delusions, hallucinations, paranoid reaction, stupor, anxiety, depression and slurred speech;

Cardiovascular: cardiac arrest, arrhythmias including malignant arrhythmias, hypotension, and tachycardia;

Respiratory: acute respiratory failure, pulmonary edema, and tachypnea;

Gastrointestinal: dysphagia;

Hematologic: leukocytosis and agranulocytosis;

Special Senses: keratitis, mydriasis, and corneal edema;

Skin and Appendages: pruritus and diaphoresis;

Miscellaneous: neuroleptic malignant syndrome (see **WARNINGS**), allergic reactions including anaphylactic reactions, edema, and fever;

Laboratory Test: elevated: CPK, BUN, serum creatinine, alkaline phosphatase, LDH, bilirubin, GGT, SGOT, and SGPT.

Overdosage

Deaths have been reported from overdose with amantadine hydrochloride. The lowest reported acute lethal dose was 1 gram. Because some patients have attempted suicide by overdosing with amantadine hydrochloride, prescriptions should be written for the smallest quantity consistent with good patient management.

Acute toxicity may be attributable to the anticholinergic effects of amantadine

hydrochloride. Drug overdose has resulted in cardiac, respiratory, renal or central nervous system toxicity. Cardiac dysfunction includes arrhythmia, tachycardia and hypertension. Pulmonary edema and respiratory distress (including adult respiratory distress syndrome -- ARDS) have been reported; renal dysfunction including increased BUN, decreased creatinine clearance and renal insufficiency can occur.

Central nervous system effects that have been reported include insomnia, anxiety, agitation, aggressive behavior, hypertonia, hyperkinesia, ataxia, gait abnormality, tremor, confusion, disorientation, depersonalization, fear, delirium, hallucinations, psychotic reactions, lethargy, somnolence and coma. Seizures may be exacerbated in patients with prior history of seizure disorders. Hyperthermia has also been observed in cases where a drug overdose has occurred.

There is no specific antidote for an overdose of amantadine hydrochloride. However, slowly administered intravenous physostigmine in 1 and 2 mg doses in an adult² at 1- to 2-hour intervals and 0.5 mg doses in a child³ at 5- to 10-minute intervals up to a maximum of 2 mg/hour have been reported to be effective in the control of central nervous system toxicity caused by amantadine hydrochloride. For acute overdosing, general supportive measures should be employed along with immediate gastric lavage or induction of emesis. Fluids should be forced, and if necessary, given intravenously. The pH of the urine has been reported to influence the excretion rate of amantadine hydrochloride. Since the excretion rate of amantadine hydrochloride increases rapidly when the urine is acidic, the administration of urine acidifying drugs may increase the elimination of the drug from the body. The blood pressure, pulse, respiration and temperature should be monitored. The patient should be observed for hyperactivity and convulsions; if required, sedation, and anticonvulsant therapy should be administered. The patient should be observed for the possible development of arrhythmias and hypotension; if required, appropriate antiarrhythmic and antihypotensive therapy should be given. Electrocardiographic monitoring may be required after ingestion, since malignant tachyarrhythmias can appear after overdose.

Care should be exercised when administering adrenergic agents, such as isoproterenol, to patients with an amantadine hydrochloride overdose, since the dopaminergic activity of amantadine hydrochloride has been reported to induce malignant arrhythmias. The blood electrolytes, urine pH and urinary output should be monitored. If there is no record of recent voiding, catheterization should be done.

Dosage and Administration

The dose of amantadine hydrochloride may need reduction in patients with congestive heart failure, peripheral edema, orthostatic hypotension, or impaired renal function (see **Dosage for Impaired Renal Function**).

Dosage for Prophylaxis and Treatment of Uncomplicated Influenza A Virus Illness:

Adult: The adult daily dosage of amantadine hydrochloride is 200 mg (four teaspoonfuls of oral solution) as a single daily dose. The daily dosage may be split into two teaspoonfuls of oral solution twice a day. If central nervous system effects develop in once-a-day dosage, a split dosage schedule may reduce such complaints.

In persons 65 years of age or older, the daily dosage of amantadine hydrochloride is 100 mg.

A 100 mg daily dose has also been shown in experimental challenge studies to be effective as prophylaxis in healthy adults who are not at high risk for influenza-related complications. However, it has not been demonstrated that a 100 mg daily dose is as

effective as a 200 mg daily dose for prophylaxis, nor has the 100 mg daily dose been studied in the treatment of acute influenza illness. In recent clinical trials, the incidence of central nervous system (CNS) side effects associated with the 100 mg daily dose was at or near the level of placebo. The 100 mg dose is recommended for persons who have demonstrated intolerance to 200 mg of amantadine hydrochloride daily because of CNS or other toxicities.

Pediatric Patients: 1 yr. to 9 yrs. of age: The total daily dose should be calculated on the basis of 2 to 4 mg/lb/day (4.4 to 8.8 mg/kg/day), but not to exceed 150 mg per day.

9 yrs. to 12 yrs. of age: The total daily dose is 200 mg given as two teaspoonfuls of oral solution twice a day. The 100 mg daily dose has not been studied in this pediatric population. Therefore, there are no data which demonstrate that this dose is as effective as or is safer than the 200 mg daily dose in this patient population.

Prophylactic dosing should be started in anticipation of an influenza A outbreak and before or after contact with individuals with influenza A virus respiratory tract illness. Amantadine hydrochloride should be continued daily for at least 10 days following a known exposure. If amantadine hydrochloride is used chemoprophylactically in conjunction with inactivated influenza A virus vaccine until protective antibody responses develop, then it should be administered for 2 to 4 weeks after the vaccine has been given. When inactivated influenza A virus vaccine is unavailable or contraindicated, amantadine hydrochloride should be administered for the duration of known influenza A in the community because of repeated and unknown exposure.

Treatment of influenza A virus illness should be started as soon as possible, preferably within 24 to 48 hours after onset of signs and symptoms, and should be continued for 24 to 48 hours after the disappearance of signs and symptoms.

Dosage for Parkinsonism:

Adult: The usual dose of amantadine hydrochloride is 100 mg twice a day when used alone. Amantadine hydrochloride has an onset of action usually within 48 hours.

The initial dose of amantadine hydrochloride is 100 mg daily for patients with serious associated medical illnesses or who are receiving high doses of other antiparkinson drugs. After one to several weeks at 100 mg once daily, the dose may be increased to 100 mg twice daily, if necessary.

Occasionally, patients whose responses are not optimal with amantadine hydrochloride at 200 mg daily may benefit from an increase up to 400 mg daily in divided doses.

However, such patients should be supervised closely by their physicians.

Patients initially deriving benefit from amantadine hydrochloride not uncommonly experience a fall-off of effectiveness after a few months. Benefit may be regained by increasing the dose to 300 mg daily. Alternatively, temporary discontinuation of amantadine hydrochloride for several weeks, followed by reinitiation of the drug, may result in regaining benefit in some patients. A decision to use other antiparkinson drugs may be necessary.

Dosage for Concomitant Therapy: Some patients who do not respond to anticholinergic antiparkinson drugs may respond to amantadine hydrochloride.

When amantadine hydrochloride or anticholinergic antiparkinson drugs are each used with marginal benefit, concomitant use may produce additional benefit.

When amantadine hydrochloride and levodopa are initiated concurrently, the patient can exhibit rapid therapeutic benefits. Amantadine hydrochloride should be held constant at 100 mg daily or twice daily while the daily dose of levodopa is gradually increased to optimal benefit.

When amantadine hydrochloride is added to optimal well-tolerated doses of levodopa, additional benefit may result, including smoothing out the fluctuations in improvement

which sometimes occur in patients on levodopa alone. Patients who require a reduction in their usual dose of levodopa because of development of side effects may possibly regain lost benefit with the addition of amantadine hydrochloride.

Dosage for Drug-Induced Extrapyrarnidal Reactions:

Adult: The usual dose of amantadine hydrochloride is 100 mg twice a day. Occasionally, patients whose responses are not optimal with amantadine hydrochloride at 200 mg daily may benefit from an increase up to 300 mg daily in divided doses.

Dosage for Impaired Renal Function:

Depending upon creatinine clearance, the following dosage adjustments are recommended:

CREATININE CLEARANCE (mL/min/1.73m ²)	AMANTADINE HYDROCHLORIDE DOSAGE
30 to 50	200 mg 1st day and 100 mg each day thereafter
15 to 29	200 mg 1st day followed by 100 mg on alternate days
<15	200 mg every 7 days

The recommended dosage for patients on hemodialysis is 200 mg every 7 days.

How Supplied

Amantadine Hydrochloride Oral Solution, USP 50 mg/5 mL is a colorless to pale yellow, raspberry-flavored oral solution available in:

10 mL unit dose cups in trays of 10 (NDC 0116-4010-10)

1 Pint (473 mL) bottles (NDC 0116-4010-16)

Store at 20° to 25°C (68° to 77°F), excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature].

KEEP TIGHTLY CLOSED

Dispense in a tight container as defined in the USP, with a child-resistant closure (as required).

Rx Only

References

1W.W. Wilson and A.H. Rajput, Amantadine-Dyazide Interaction, Can. Med. Assoc. J. 129:974-975, 1983.

2D.F. Casey, N. Engl. J. Med. 298:516, 1978.

3C.D. Berkowitz, J. Pediatr. 95:144, 1979.

Manufactured by:

Xttrium Laboratories, Inc.
1200 E. Business Center Dr.
Mount Prospect, IL 60056

4010AMANINSB REV. 06-25

Amantadine 16oz Label

Each 5 mL (1 teaspoonful) contains: Amantadine Hydrochloride, USP 50 mg USUAL DOSAGE: See accompanying package insert for complete prescribing information. WARNINGS: KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN. In case of accidental overdose, seek professional assistance or contact a poison control center immediately. Store at 20° to 25°C (68° to 77°F), excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature]. KEEP TIGHTLY CLOSED Dispense in a tight container as defined in the USP, with a child-resistant closure (as required). Manufactured by: Xttrium Laboratories, Inc. 1200 E. Business Center Dr. Mount Prospect, IL 60056 401016OZLBC REV. 06-25  N 0 01164 01016 4	 NDC 0116-4010-16 AMANTADINE HYDROCHLORIDE ORAL SOLUTION, USP 50 mg/5 mL DO NOT USE IF TAMPER-EVIDENT SEAL IS BROKEN OR MISSING. Rx Only NET: 1 Pint (473 mL) 
--	--


Amantadine Hydrochloride Oral Solution, USP

50 mg/5 mL

Rx only

DESCRIPTION

Amantadine hydrochloride, USP is designated generically as amantadine hydrochloride and chemically as 1-adamantanamine hydrochloride.



• HCl

$C_{10}H_{17}N \bullet HCl$

M.W. 187.71

Amantadine hydrochloride is a stable white or nearly white crystalline powder, freely soluble in water and soluble in alcohol and in chloroform.

Amantadine hydrochloride has pharmacological actions as both an anti-Parkinson and an antiviral drug.

Amantadine Hydrochloride Oral Solution, USP contains 50 mg of amantadine hydrochloride per 5 mL and has the following inactive ingredients: anhydrous citric acid, artificial raspberry flavor, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, sodium citrate dihydrate, and sorbitol solution.

CLINICAL PHARMACOLOGY

Pharmacodynamics

Mechanism of Action: Antiviral: The mechanism by which amantadine hydrochloride exerts its antiviral activity is not clearly understood. It appears to mainly prevent the release of infectious viral nucleic acid into the host cell by interfering with the function of the transmembrane domain of the viral M2 protein. In certain cases, amantadine hydrochloride is also known to prevent virus assembly during virus replication. It does not appear to interfere with the immunogenicity of inactivated influenza A virus vaccine.

Antiviral Activity: Amantadine hydrochloride inhibits the replication of influenza A virus isolates from each of the subtypes, i.e., H1N1, H2N2 and H3N2. It has very little or no activity against influenza B virus isolates. A quantitative relationship between the *in vitro* susceptibility of influenza A virus to amantadine hydrochloride and the clinical response to therapy has not been established in man. Sensitivity test results, expressed as the concentration of amantadine hydrochloride required to inhibit by 50% the growth of virus (ED_{50}) in tissue culture vary greatly (from 0.1 $\mu\text{g/mL}$ to 25.0 $\mu\text{g/mL}$) depending upon the assay protocol used, size of virus inoculum, isolates of influenza A virus strains tested, and the cell type used. Host cells in tissue culture readily tolerated amantadine hydrochloride up to a concentration of 100 $\mu\text{g/mL}$.

Mechanism of Action: Parkinson's Disease: The mechanism of action of amantadine hydrochloride in the treatment of Parkinson's disease and drug-induced extrapyramidal reactions is not known. Data from earlier animal studies suggest that amantadine hydrochloride may have direct and indirect effects on dopamine neurons. More recent studies have demonstrated that amantadine hydrochloride is a weak, non-competitive NMDA receptor antagonist ($K_i = 10\mu\text{M}$). Although amantadine hydrochloride has not been shown to possess direct anticholinergic activity in animal studies, clinically, it exhibits anticholinergic-like side effects such as dry mouth, urinary retention, and constipation.

Pharmacokinetics

Amantadine hydrochloride is well absorbed orally. Maximum plasma concentrations are directly related to dose for doses up to 200 mg/day. Doses above 200 mg/day may result in a greater than proportional increase in maximum plasma concentrations. It is primarily excreted unchanged in the urine by glomerular filtration and tubular secretion. Eight metabolites of amantadine hydrochloride have been identified in human urine. One metabolite, an N-acetylated compound, was quantified in human urine and accounted for 5 to 15% of the administered dose. Plasma acetylamantadine accounted for up to 80% of the concurrent amantadine hydrochloride plasma concentration in 5 of 12 healthy volunteers following the ingestion of a 200 mg dose of amantadine hydrochloride. Acetylamantadine was not detected in the plasma of the remaining seven volunteers. The contribution of this metabolite to efficacy or toxicity is not known.

There appears to be a relationship between plasma amantadine hydrochloride concentrations and toxicity. As concentration increases, toxicity seems to be more prevalent, however, absolute values of amantadine hydrochloride concentrations associated with adverse effects have not been fully defined.

Amantadine hydrochloride pharmacokinetics were determined in 24 normal adult male volunteers after the oral administration of a single amantadine hydrochloride 100 mg soft gel capsule. The mean $\pm$ SD maximum plasma concentration was $0.22 \pm 0.03 \mu\text{g/mL}$ (range: 0.18 to 0.32 $\mu\text{g/mL}$). The time to peak concentration was 3.3 ± 1.5 hours (range: 1.5 to 8.0 hours). The apparent oral clearance was $0.28 \pm 0.11 \text{ L/hr/kg}$ (range: 0.14 to 0.62 L/hr/kg). The half-life was 17 ± 4 hours (range: 10 to 25 hours). Across other studies, amantadine hydrochloride plasma half-life has averaged 16 ± 6 hours (range: 9 to 31 hours) in 19 healthy volunteers.

After oral administration of a single dose of 100 mg amantadine hydrochloride in a syrup formulation to five healthy volunteers, the mean $\pm$ SD maximum plasma concentration C_{max} was $0.24 \pm 0.04 \mu\text{g/mL}$ and ranged from 0.18 to 0.28 $\mu\text{g/mL}$. After 15 days of amantadine hydrochloride 100 mg b.i.d., the C_{max} was $0.47 \pm 0.11 \mu\text{g/mL}$ in four of the five volunteers. Across studies, the time to C_{max} (T_{max}) averaged about 2 to 4 hours.

Plasma amantadine hydrochloride clearance ranged from 0.2 to 0.3 L/hr/kg after the administration of 5 mg to 25 mg intravenous doses of amantadine hydrochloride to 15 healthy volunteers.

In six healthy volunteers, the ratio of amantadine hydrochloride renal clearance to apparent oral plasma clearance was 0.79 ± 0.17 (mean $\pm$ SD).

The volume of distribution determined after the intravenous administration of amantadine hydrochloride to 15 healthy subjects was 3 to 8 L/kg, suggesting tissue binding. Amantadine hydrochloride, after single oral 200 mg doses to 6 healthy young subjects and to 6 healthy elderly subjects has been found in nasal mucus at mean $\pm$ SD concentrations of 0.15 ± 0.16 , 0.28 ± 0.26 , and 0.39 ± 0.34 $\mu\text{g/g}$ at 1, 4, and 8 hours after dosing, respectively. These concentrations represented $31 \pm 33\%$, $59 \pm 61\%$, and $95 \pm 86\%$ of the corresponding plasma amantadine hydrochloride concentrations. Amantadine hydrochloride is approximately 67% bound to plasma proteins over a concentration range of 0.1 to 2.0 $\mu\text{g/mL}$. Following the administration of amantadine hydrochloride 100 mg as a single dose, the mean $\pm$ SD red blood cell to plasma ratio ranged from 2.7 ± 0.5 in 6 healthy subjects to 1.4 ± 0.2 in 8 patients with renal insufficiency.

The apparent oral plasma clearance of amantadine hydrochloride is reduced and the plasma half-life and plasma concentrations are increased in healthy elderly individuals age 60 and older. After single dose administration of 25 to 75 mg to 7 healthy, elderly male volunteers, the apparent plasma clearance of amantadine hydrochloride was 0.10 ± 0.04 L/hr/kg (range 0.06 to 0.17 L/hr/kg) and the half-life was 29 ± 7 hours (range 20 to 41 hours). Whether these changes are due to decline in renal function or other age related factors is not known.

In a study of young healthy subjects ($n=20$), mean renal clearance of amantadine hydrochloride, normalized for body mass index, was 1.5 fold higher in males compared to females ($p<0.032$).

Compared with otherwise healthy adult individuals, the clearance of amantadine hydrochloride is significantly reduced in adult patients with renal insufficiency. The elimination half-life increases two to three fold or greater when creatinine clearance is less than 40 mL/min/1.73 m² and averages eight days in patients on chronic maintenance hemodialysis. Amantadine hydrochloride is removed in negligible amounts by hemodialysis.

The pH of the urine has been reported to influence the excretion rate of amantadine hydrochloride. Since the excretion rate of amantadine hydrochloride increases rapidly when the urine is acidic, the administration of urine acidifying drugs may increase the elimination of the drug from the body.

INDICATIONS AND USAGE

Amantadine Hydrochloride Oral Solution, USP is indicated for the prophylaxis and treatment of signs and symptoms of infection caused by various strains of influenza A virus. Amantadine hydrochloride is also indicated in the treatment of parkinsonism and drug-induced extrapyramidal reactions.

Influenza A Prophylaxis: Amantadine hydrochloride is indicated for chemoprophylaxis against signs and symptoms of influenza A virus infection. Because amantadine hydrochloride does not completely prevent the host immune response to influenza A infection, individuals who take this drug may still develop immune responses to natural disease or vaccination and may be protected when later exposed to antigenically related viruses. Following vaccination during an influenza A outbreak, amantadine hydrochloride prophylaxis should be considered for the 2- to 4-week time period required to develop an antibody response.

Influenza A Treatment: Amantadine hydrochloride is also indicated in the treatment of uncomplicated respiratory tract illness caused by influenza A virus strains especially when administered early in the course of illness. There are no well-controlled clinical studies demonstrating that treatment with amantadine hydrochloride will avoid the development of influenza A virus pneumonitis or other complications in high risk patients.

There is no clinical evidence indicating that amantadine hydrochloride is effective in the prophylaxis or treatment of viral respiratory tract illnesses other than those caused by influenza A virus strains.

The following points should be considered before initiating treatment or prophylaxis with amantadine hydrochloride:

- Amantadine hydrochloride is not a substitute for early vaccination on an annual basis as recommended by the Centers for Disease Control and Prevention Advisory Committee on Immunization Practices.
- Influenza viruses change over time. Emergence of resistance mutations could decrease drug effectiveness. Other factors (for example, changes in viral virulence) might also diminish clinical benefit of antiviral drugs. Prescribers should consider available information on influenza drug susceptibility patterns and treatment effects when deciding whether to use amantadine hydrochloride.

Parkinson's Disease/Syndrome: Amantadine hydrochloride is indicated in the treatment of idiopathic Parkinson's disease (Paralysis Agitans), postencephalitic parkinsonism, and symptomatic parkinsonism which may follow injury to the nervous system by carbon monoxide intoxication. It is indicated in those elderly patients believed to develop parkinsonism in association with cerebral arteriosclerosis. In the treatment of Parkinson's disease, amantadine hydrochloride is less effective than levodopa, (-)-3-(3,4-dihydroxyphenyl)-L-alanine, and its efficacy in comparison with the anticholinergic antiparkinson drugs has not yet been established.

Drug-Induced Extrapyramidal Reactions:

Amantadine hydrochloride is indicated in the treatment of drug-induced extrapyramidal reactions. Although anticholinergic-type side effects have been noted with amantadine hydrochloride when used in patients with drug-induced extrapyramidal reactions, there is a lower incidence of these side effects than that observed with the anticholinergic antiparkinson drugs.

CONTRAINDICATIONS

Amantadine hydrochloride is contraindicated in patients with known hypersensitivity to amantadine hydrochloride or to any of the other ingredients in Amantadine Hydrochloride Oral Solution, USP.

WARNINGS

Deaths: Deaths have been reported from overdose with amantadine hydrochloride. The lowest reported acute lethal dose was 1 gram. Acute toxicity may be attributable to the anticholinergic effects of amantadine hydrochloride. Drug overdose has resulted in cardiac, respiratory, renal or central nervous system toxicity. Cardiac dysfunction includes arrhythmia, tachycardia and hypertension (see **OVERDOSAGE**).

Deaths due to drug accumulation (overdose) have been reported in patients with renal impairment, who were prescribed higher than recommended doses of amantadine hydrochloride for their level of renal function (see **DOSAGE AND ADMINISTRATION; Dosage of Impaired Renal Function** and **OVERDOSAGE**).

Suicide Attempts: Suicide attempts, some of which have been fatal, have been reported in patients treated with amantadine hydrochloride, many of whom received short courses for influenza treatment or prophylaxis. The incidence of suicide attempts is not known and the pathophysiologic mechanism is not understood. Suicide attempts and suicidal ideation have been reported in patients with and without prior history of psychiatric illness. Amantadine hydrochloride can exacerbate mental problems in patients with a history of psychiatric disorders or substance abuse. Patients who attempt suicide may exhibit abnormal mental states which include disorientation, confusion, depression, personality changes, agitation, aggressive behavior, hallucinations, paranoia, other psychotic reactions, and somnolence or insomnia. Because of the possibility of serious adverse effects, caution should be observed when prescribing amantadine hydrochloride to patients being treated with drugs having CNS effects, or for whom the potential risks outweigh the benefit of treatment.

CNS Effects: Patients with a history of epilepsy or other "seizures" should be observed closely for possible increased seizure activity.

Patients receiving amantadine hydrochloride who note central nervous system effects or blurring of vision should be cautioned against driving or working in situations where alertness and adequate motor coordination are important.

Other: Patients with a history of congestive heart failure or peripheral edema should be followed closely as there are patients who developed congestive heart failure while receiving amantadine hydrochloride.

Patients with Parkinson's disease improving on amantadine hydrochloride should resume normal activities gradually and cautiously, consistent with other medical considerations, such as the presence of osteoporosis or phlebothrombosis.

Because amantadine hydrochloride has anticholinergic effects and may cause mydriasis, it should not be given to patients with untreated angle closure glaucoma.

Corneal Edema: Corneal edema has been reported in patients taking amantadine. Symptoms include sudden onset of blurry vision, or progressive vision loss, with or without eye pain. Corneal involvement is usually bilateral. Onset can occur from a few weeks to several years after starting amantadine. Resolution of symptoms typically begins within weeks of amantadine cessation. However, corneal grafts have been required in some patients when the condition is not recognized. Permanent damage can occur if amantadine is continued. Ask patients if their vision has changed and obtain ophthalmologic examinations to rule out corneal edema should vision changes occur after initiation of therapy with amantadine. If corneal edema occurs, taper and discontinue amantadine [see **DOSAGE AND ADMINISTRATION**].

PRECAUTIONS

Amantadine hydrochloride should not be discontinued abruptly in patients with Parkinson's disease since a few patients have experienced a parkinsonian crisis, i.e., a sudden marked clinical deterioration, when this medication was suddenly stopped. The dose of anticholinergic drugs or of amantadine hydrochloride should be reduced if atropine-like effects appear when these drugs are used concurrently. Abrupt discontinuation may also precipitate delirium, agitation, delusions, hallucinations, paranoid reaction, stupor, anxiety, depression and slurred speech.

Neuroleptic Malignant Syndrome (NMS): Sporadic cases of possible Neuroleptic Malignant Syndrome (NMS) have been reported in association with dose reduction or withdrawal of amantadine hydrochloride therapy. Therefore, patients should be observed carefully when the dosage of amantadine hydrochloride is reduced abruptly or discontinued, especially if the patient is receiving neuroleptics.

NMS is an uncommon but life-threatening syndrome characterized by fever or hyperthermia; neurologic findings including muscle rigidity, involuntary movements, altered consciousness; mental status changes; other disturbances such as autonomic dysfunction, tachycardia, tachypnea, hyper- or hypotension; laboratory findings such as creatine phosphokinase elevation, leukocytosis, myoglobinuria, and increased serum myoglobin.

The early diagnosis of this condition is important for the appropriate management of these patients. Considering NMS as a possible diagnosis and ruling out other acute illnesses (e.g., pneumonia, systemic infection, etc.) is essential. This may be especially complex if the clinical presentation includes both serious medical illness and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever and primary central nervous system (CNS) pathology.

The management of NMS should include: 1) intensive symptomatic treatment and medical monitoring, and 2) treatment of any concomitant serious medical problems for which specific treatments are available. Dopamine agonists, such as bromocriptine, and muscle relaxants, such as dantrolene are often used in the treatment of NMS, however, their effectiveness has not been demonstrated in controlled studies.

Renal disease: Because amantadine hydrochloride is mainly excreted in the urine, it accumulates in the plasma and in the body when renal function declines. Thus, the dose of amantadine hydrochloride should be reduced in patients with renal impairment and in individuals who are 65 years of age or older (see **DOSAGE AND ADMINISTRATION**; Dosage for Impaired Renal Function).

Liver disease: Care should be exercised when administering amantadine hydrochloride to patients with liver disease. Rare instances of reversible elevation of liver enzymes have been reported in patients receiving amantadine hydrochloride, though a specific relationship between the drug and such changes has not been established.

Impulse Control/Compulsive Behaviors:

Postmarketing reports suggest that patients treated with anti-Parkinson medications can experience intense urges to gamble, increased sexual urges, intense urges to spend money uncontrollably, and other intense urges. Patients may be unable to control these urges while taking one or more of the medications that are generally used for the treatment of Parkinson's disease and that increase central dopaminergic tone, including Amantadine hydrochloride. In some cases, although not all, these urges were reported to have stopped when the dose was reduced or the medication was discontinued. Because patients may not recognize these behaviors as abnormal it is important for prescribers to specifically ask patients or their caregivers about the development of new or increased gambling urges, sexual urges, uncontrolled spending or other urges while being treated with Amantadine hydrochloride. Physicians should consider dose reduction or stopping the medication if a patient develops such urges while taking amantadine hydrochloride.

Melanoma:

Epidemiological studies have shown that patients with Parkinson's disease have a higher risk (2- to approximately 6-fold higher) of developing melanoma than the general population. Whether the increased risk observed was due to Parkinson's disease or other factors, such as drugs used to treat Parkinson's disease, is unclear.

For the reasons stated above, patients and providers are advised to monitor for melanomas frequently and on a regular basis when using amantadine hydrochloride for any indication. Ideally, periodic skin examinations should be performed by appropriately qualified individuals (e.g., dermatologists).

Other: The dose of amantadine hydrochloride may need careful adjustment in patients with congestive heart failure, peripheral edema, or orthostatic hypotension. Care should be exercised when administering amantadine hydrochloride to patients with a history of recurrent eczematoid rash, or to patients with psychosis or severe psychoneurosis not controlled by chemotherapeutic agents.

Serious bacterial infections may begin with influenza-like symptoms or may coexist with or occur as complications during the course of influenza. Amantadine hydrochloride has not been shown to prevent such complications.

Information for Patients: Patients should be advised of the following information:

Blurry vision, eye pain, loss of vision, and/or impaired mental acuity may occur.

Gradually increase physical activity as the symptoms of Parkinson's disease improve.

Avoid excessive alcohol usage, since it may increase the potential for CNS effects such as dizziness, confusion, lightheadedness and orthostatic hypotension.

Avoid getting up suddenly from a sitting or lying position. If dizziness or lightheadedness occurs, notify physician.

Notify physician if mood/mental changes, swelling of extremities, difficulty urinating and/or shortness of breath occur.

Do not take more medication than prescribed because of the risk of overdose. If there is no improvement in a few days, or if medication appears less effective after a few weeks, discuss with a physician.

Consult physician before discontinuing medication. Seek medical attention immediately if it is suspected that an overdose of medication has been taken.

Drug Interactions: Careful observation is required when amantadine hydrochloride is administered concurrently with central nervous system stimulants.

Agents with anticholinergic properties may potentiate the anticholinergic-like side effects of amantadine hydrochloride.

Coadministration of thioridazine has been reported to worsen the tremor in elderly patients with Parkinson's disease, however, it is not known if other phenothiazines produce a similar response.

Coadministration of triamterene/hydrochlorothiazide resulted in a higher plasma amantadine hydrochloride concentration in a 61-year-old man receiving amantadine hydrochloride 100 mg TID for Parkinson's disease.¹ It is not known which of the components of triamterene/hydrochlorothiazide contributed to the observation or if related drugs produce a similar response.

Coadministration of quinine or quinidine with amantadine hydrochloride was shown to reduce the renal clearance of amantadine hydrochloride by about 30%.

The concurrent use of amantadine hydrochloride with live attenuated influenza vaccine (LAV) intranasal has not been evaluated. However, because of the potential for interference between these products, LAV should not be administered within 2 weeks before or 48 hours after administration of amantadine hydrochloride, unless medically indicated. The concern about possible interference arises from the potential for antiviral drugs to inhibit replication of live vaccine virus. Trivalent inactivated influenza vaccine can be administered at any time relative to use of amantadine hydrochloride.

Carcinogenesis, Mutagenesis: Long-term *in vivo* animal studies designed to evaluate the carcinogenic potential of amantadine hydrochloride have not been performed. In several *in vitro* assays for gene mutation, amantadine hydrochloride did not increase the number of spontaneously observed mutations in four strains of *Salmonella typhimurium* (Ames Test) or in a mammalian cell line (Chinese Hamster Ovary cells) when incubations were performed either with or without a liver metabolic activation extract. Further, there was no evidence of chromosome damage observed in an *in vitro* test using freshly derived and stimulated human peripheral blood lymphocytes (with and without metabolic activation) or in an *in vivo* mouse bone marrow micronucleus test (140 to 550 mg/kg; estimated human equivalent doses of 11.7 to 45.8 mg/kg based on body surface area conversion).

Impairment of Fertility: The effect of amantadine hydrochloride on fertility has not been adequately tested, that is, in a study conducted under Good Laboratory Practice (GLP) and according to current recommended methodology. In a three litter, non-GLP, reproduction study in rats, amantadine hydrochloride at a dose of 32 mg/kg/day (equal to the maximum recommended human dose on a mg/m² basis) administered to both males and females slightly impaired fertility. There were no effects on fertility at a dose level of 10 mg/kg/day (or 0.3 times the maximum recommended human dose on a mg/m² basis); intermediate doses were not tested.

Failed fertility has been reported during human in vitro fertilization (IVF) when the sperm donor ingested amantadine hydrochloride 2 weeks prior to, and during the IVF cycle.

Pregnancy:

Teratogenic Effects: The effect of amantadine hydrochloride on embryofetal and peri-postnatal development has not been adequately tested, that is, in studies conducted under Good Laboratory Practice (GLP) and according to current recommended methodology. However, in two non-GLP studies in rats in which females were dosed from 5 days prior to mating to Day 6 of gestation or on Days 7 to 14 of gestation, amantadine hydrochloride produced increases in embryonic death at an oral dose of 100 mg/kg (or 3 times the maximum recommended human dose on a mg/m² basis). In the non-GLP rat study in which females were dosed on Days 7 to 14 of gestation, there was a marked increase in severe visceral and skeletal malformations at oral doses of 50 and 100 mg/kg (or 1.5 and 3 times, respectively, the maximum recommended human dose on a mg/m² basis). The no-effect dose for teratogenicity was 37 mg/kg (equal to the maximum recommended human dose on a mg/m² basis). The safety margins reported may not accurately reflect the risk considering the questionable quality of the study on which they are based. There are no adequate and well-controlled studies in pregnant women. Human data regarding teratogenicity after maternal use of amantadine hydrochloride is scarce. Tetralogy of Fallot and tibial hemimelia (normal karyotype) occurred in an infant exposed to amantadine hydrochloride during the first trimester of pregnancy (100 mg P.O. for 7 days during the 6th and 7th week of gestation). Cardiovascular maldevelopment (single ventricle with pulmonary atresia) was associated with maternal exposure to amantadine hydrochloride (100 mg/d) administered during the first 2 weeks of pregnancy. Amantadine hydrochloride should be used during pregnancy only if the potential benefit justifies the potential risk to the embryo or fetus.

Nursing Mothers: Amantadine hydrochloride is excreted in human milk. Use is not recommended in nursing mothers.

Pediatric Use: The safety and efficacy of amantadine hydrochloride in newborn infants and infants below the age of 1 year have not been established.

Usage in the Elderly: Because amantadine hydrochloride is primarily excreted in the urine, it accumulates in the plasma and in the body when renal function declines. Thus, the dose of amantadine hydrochloride should be reduced in patients with renal impairment and in individuals who are 65 years of age or older. The dose of amantadine hydrochloride may need reduction in patients with congestive heart failure, peripheral edema, or orthostatic hypotension (see **DOSAGE AND ADMINISTRATION**).

ADVERSE REACTIONS

The adverse reactions reported most frequently at the recommended dose of amantadine hydrochloride (5 to 10%) are: nausea, dizziness (lightheadedness), and insomnia.

Less frequently (1 to 5%) reported adverse reactions are:

depression, anxiety and irritability, hallucinations, confusion, anorexia, dry mouth, constipation, ataxia, livedo reticularis, peripheral edema, orthostatic hypotension, headache, somnolence, nervousness, dream abnormality, agitation, dry nose, diarrhea and fatigue.

Infrequently (0.1 to 1%) occurring adverse reactions are: congestive heart failure, psychosis, urinary retention, dyspnea, skin rash, vomiting, weakness, slurred speech, euphoria, thinking abnormality, amnesia, hyperkinesia, hypertension, decreased libido, and visual disturbance, including punctate subepithelial or other corneal opacity, corneal edema, decreased visual acuity, sensitivity to light, and optic nerve palsy.

Rare (less than 0.1%) occurring adverse reactions are:

instances of convulsion, leukopenia, neutropenia, eczematoid dermatitis, oculogyric episodes, suicidal attempt, suicide, and suicidal ideation (see **WARNINGS**).

Other adverse reactions reported during postmarketing experience with amantadine hydrochloride usage include:

Nervous System/Psychiatric: coma, stupor, delirium, hypokinesia, hypertonia, delusions, aggressive behavior, paranoid reaction, manic reaction, involuntary muscle contractions, gait abnormalities, paresthesia, EEG changes, and tremor. Abrupt discontinuation may also precipitate delirium, agitation, delusions, hallucinations, paranoid reaction, stupor, anxiety, depression and slurred speech;

Cardiovascular: cardiac arrest, arrhythmias including malignant arrhythmias, hypotension, and tachycardia;

Respiratory: acute respiratory failure, pulmonary edema, and tachypnea;

Gastrointestinal: dysphagia;

Hematologic: leukocytosis and agranulocytosis;

Special Senses: keratitis, mydriasis, and corneal edema;

Skin and Appendages: pruritus and diaphoresis;

Miscellaneous: neuroleptic malignant syndrome (see **WARNINGS**), allergic reactions including anaphylactic reactions, edema, and fever;

Laboratory Test: elevated: CPK, BUN, serum creatinine, alkaline phosphatase, LDH, bilirubin, GGT, SGOT, and SGPT.

OVERDOSAGE

Deaths have been reported from overdose with amantadine hydrochloride. The lowest reported acute lethal dose was 1 gram. Because some patients have attempted suicide by overdosing with amantadine hydrochloride, prescriptions should be written for the smallest quantity consistent with good patient management.

Acute toxicity may be attributable to the anticholinergic effects of amantadine hydrochloride. Drug overdose has resulted in cardiac, respiratory, renal or central nervous system toxicity. Cardiac dysfunction includes arrhythmia, tachycardia and hypertension. Pulmonary edema and respiratory distress (including adult respiratory distress syndrome — ARDS) have been reported; renal dysfunction including increased BUN, decreased creatinine clearance and renal insufficiency can occur.

Central nervous system effects that have been reported include insomnia, anxiety, agitation, aggressive behavior, hypertonia, hyperkinesia, ataxia, gait abnormality, tremor, confusion, disorientation, depersonalization, fear, delirium, hallucinations, psychotic reactions, lethargy, somnolence and coma. Seizures may be exacerbated in patients with prior history of seizure disorders. Hyperthermia has also been observed in cases where a drug overdose has occurred.

There is no specific antidote for an overdose of amantadine hydrochloride. However, slowly administered intravenous physostigmine in 1 and 2 mg doses in an adult² at 1- to 2-hour intervals and 0.5 mg doses in a child³ at 5- to 10-minute intervals up to a maximum of 2 mg/hour have been reported to be effective in the control of central nervous system toxicity caused by amantadine hydrochloride. For acute overdosing, general supportive measures should be employed along with immediate gastric lavage or induction of emesis. Fluids should be forced, and if necessary, given intravenously. The pH of the urine has been reported to influence the excretion rate of amantadine hydrochloride. Since the excretion rate of amantadine hydrochloride increases rapidly when the urine is acidic, the administration of urine acidifying drugs may increase the elimination of the drug from the body. The blood pressure, pulse, respiration and temperature should be monitored. The patient should be observed for hyperactivity and convulsions; if required, sedation, and anticonvulsant therapy should be administered. The patient should be observed for the possible development of arrhythmias and hypotension; if required, appropriate antiarrhythmic and antihypotensive therapy should be given. Electrocardiographic monitoring may be required after ingestion, since malignant tachyarrhythmias can appear after overdose.

Care should be exercised when administering adrenergic agents, such as isoproterenol, to patients with an amantadine hydrochloride overdose, since the dopaminergic activity of amantadine hydrochloride has been reported to induce malignant arrhythmias.

The blood electrolytes, urine pH and urinary output should be monitored. If there is no record of recent voiding, catheterization should be done.

DOSAGE AND ADMINISTRATION

The dose of amantadine hydrochloride may need reduction in patients with congestive heart failure, peripheral edema, orthostatic hypotension, or impaired renal function (see **Dosage for Impaired Renal Function**).

Dosage for Prophylaxis and Treatment of Uncomplicated Influenza A Virus Illness:

Adult: The adult daily dosage of amantadine hydrochloride is 200 mg (four teaspoonfuls of oral solution) as a single daily dose. The daily dosage may be split into two teaspoonfuls of oral solution twice a day. If central nervous system effects develop in once-a-day dosage, a split dosage schedule may reduce such complaints.

In persons 65 years of age or older, the daily dosage of amantadine hydrochloride is 100 mg. A 100 mg daily dose has also been shown in experimental challenge studies to be effective as prophylaxis in healthy adults who are not at high risk for influenza-related complications. However, it has not been demonstrated that a 100 mg daily dose is as effective as a 200 mg daily dose for prophylaxis, nor has the 100 mg daily dose been studied in the treatment of acute influenza illness. In recent clinical trials, the incidence of central nervous system (CNS) side effects associated with the 100 mg daily dose was at or near the level of placebo. The 100 mg dose is recommended for persons who have demonstrated intolerance to 200 mg of amantadine hydrochloride daily because of CNS or other toxicities.

Pediatric Patients: 1 yr. to 9 yrs. of age: The total daily dose should be calculated on the basis of 2 to 4 mg/lb/day (4.4 to 8.8 mg/kg/day), but not to exceed 150 mg per day.

9 yrs. to 12 yrs. of age: The total daily dose is 200 mg given as two teaspoonfuls of oral solution twice a day. The 100 mg daily dose has not been studied in this pediatric population. Therefore, there are no data which demonstrate that this dose is as effective as or is safer than the 200 mg daily dose in this patient population.

Prophylactic dosing should be started in anticipation of an influenza A outbreak and before or after contact with individuals with influenza A virus respiratory tract illness.

Amantadine hydrochloride should be continued daily for at least 10 days following a known exposure. If amantadine hydrochloride is used chemoprophylactically in conjunction with inactivated influenza A virus vaccine until protective antibody responses develop, then it should be administered for 2 to 4 weeks after the vaccine has been given. When inactivated influenza A virus vaccine is unavailable or contraindicated, amantadine hydrochloride should be administered for the duration of known influenza A in the community because of repeated and unknown exposure.

Treatment of influenza A virus illness should be started as soon as possible, preferably within 24 to 48 hours after onset of signs and symptoms, and should be continued for 24 to 48 hours after the disappearance of signs and symptoms.

Dosage for Parkinsonism:

Adult: The usual dose of amantadine hydrochloride is 100 mg twice a day when used alone. Amantadine hydrochloride has an onset of action usually within 48 hours.

The initial dose of amantadine hydrochloride is 100 mg daily for patients with serious associated medical illnesses or who are receiving high doses of other antiparkinson drugs. After one to several weeks at 100 mg once daily, the dose may be increased to 100 mg twice daily, if necessary.

Occasionally, patients whose responses are not optimal with amantadine hydrochloride at 200 mg daily may benefit from an increase up to 400 mg daily in divided doses. However, such patients should be supervised closely by their physicians.

Patients initially deriving benefit from amantadine hydrochloride not uncommonly experience a fall-off of effectiveness after a few months. Benefit may be regained by increasing the dose to 300 mg daily. Alternatively, temporary discontinuation of amantadine hydrochloride for several weeks, followed by reinitiation of the drug, may result in regaining benefit in some patients. A decision to use other antiparkinson drugs may be necessary.

Dosage for Concomitant Therapy: Some patients who do not respond to anticholinergic antiparkinson drugs may respond to amantadine hydrochloride.

When amantadine hydrochloride or anticholinergic antiparkinson drugs are each used with marginal benefit, concomitant use may produce additional benefit.

When amantadine hydrochloride and levodopa are initiated concurrently, the patient can exhibit rapid therapeutic benefits. Amantadine hydrochloride should be held constant at 100 mg daily or twice daily while the daily dose of levodopa is gradually increased to optimal benefit.

When amantadine hydrochloride is added to optimal well-tolerated doses of levodopa, additional benefit may result, including smoothing out the fluctuations in improvement which sometimes occur in patients on levodopa alone. Patients who require a reduction in their usual dose of levodopa because of development of side effects may possibly regain lost benefit with the addition of amantadine hydrochloride.

Dosage for Drug-Induced Extrapyrimal Reactions:

Adult: The usual dose of amantadine hydrochloride is 100 mg twice a day. Occasionally, patients whose responses are not optimal with amantadine hydrochloride at 200 mg daily may benefit from an increase up to 300 mg daily in divided doses.

Dosage for Impaired Renal Function:

Depending upon creatinine clearance, the following dosage adjustments are recommended:

CREATININE CLEARANCE (mL/min/1.73m ²)	AMANTADINE HYDROCHLORIDE DOSAGE
30 to 50	200 mg 1st day and 100 mg each day thereafter
15 to 29	200 mg 1st day followed by 100 mg on alternate days
<15	200 mg every 7 days

The recommended dosage for patients on hemodialysis is 200 mg every 7 days.

HOW SUPPLIED

Amantadine Hydrochloride Oral Solution, USP 50 mg/5 mL is a colorless to pale yellow, raspberry-flavored oral solution available in:

10 mL unit dose cups in trays of 10 (NDC 0116-4010-10)

1 Pint (473 mL) bottles (NDC 0116-4010-16)

**Store at 20° to 25°C (68° to 77°F), excursions permitted to 15° to 30°C (59° to 86°F).
[See USP Controlled Room Temperature].**

KEEP TIGHTLY CLOSED

Dispense in a tight container as defined in the USP, with a child-resistant closure (as required).

Rx Only

REFERENCES

¹W.W. Wilson and A.H. Rajput, Amantadine-Dyazide Interaction, *Can. Med. Assoc. J.* 129:974-975, 1983.

²D.F. Casey, *N. Engl. J. Med.* 298:516, 1978.

³C.D. Berkowitz, *J. Pediatr.* 95:144, 1979.

Manufactured by:

Xttrium Laboratories, Inc.

1200 E. Business Center Dr.

Mount Prospect, IL 60056

4010160ZLBLC

REV. 06-25

Each 5 mL (1 teaspoonful) contains:

Amantadine Hydrochloride, USP 50 mg

USUAL DOSAGE: See accompanying package insert for complete prescribing information.

WARNINGS: KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

In case of accidental overdose, seek professional assistance or contact a poison control center immediately.

Store at 20° to 25°C (68° to 77°F), excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature]. KEEP TIGHTLY CLOSED

Dispense in a tight container as defined in the USP, with a child-resistant closure (as required).

Manufactured by:

Xttrium Laboratories, Inc.

1200 E. Business Center Dr.

Mount Prospect, IL 60056

4010160ZLBLC

REV. 06-25



Amantadine 10mL Lid

UNIT DOSE
Delivers 10 mL
NDC 0116-4010-10
Amantadine Hydrochloride
Oral Solution USP,
100 mg per 10 mL



301164010103

4010AMANI01IDA

Rx Only
FOR INSTITUTIONAL USE ONLY
Xttrium Laboratories, Inc.
Mount Prospect
IL 60056
Usual Dosage: See Insert

xl[®]

Amantadine Insert

Thus, the dose of amantadine hydrochloride should be reduced in patients with renal impairment and in individuals who are 65 years of age or older (see DOSAGE AND ADMINISTRATION, Dosage for Impaired Renal Function). Urinary Diseases: Care should be exercised when administering amantadine hydrochloride to patients with urinary diseases. Past instances of reversible elevation of liver enzymes have been reported in patients receiving amantadine hydrochloride, though a specific relationship between this drug and such changes has not been established. Impaired Control/Compulsive Behaviors: Postmarketing reports suggest that patients treated with anti-Parkinson medications can experience intense urges to gamble, increased sexual urges, intense urges to spend money uncontrollably, and/or other intense urges. Patients may be unable to control these urges while taking one or more of the medications that are generally used for the treatment of Parkinson's disease and that increase central dopaminergic tone, including amantadine hydrochloride. In some cases, although not all, these urges were reported to have stopped when the dose was reduced or the medication was discontinued. Because patients may not recognize these behaviors as abnormal if they have been taking amantadine hydrochloride, patients or their caregivers about the development of new or increased gambling urges, sexual urges, uncontrolled spending or other urges while taking treatment with Amantadine hydrochloride. Physicians should consider dose reduction or stopping the medication if a patient develops such urges while taking amantadine hydrochloride. Melaninemia: Epidemiological studies have shown that patients with Parkinson's disease have a higher risk (2- to approximately 6-fold higher) of developing melanoma than the general population. Whether the increase effect observed and due to Parkinson's Disease or other factors, such as drugs used to treat Parkinson's disease, is unclear. For the reasons stated above, patients and providers are advised to monitor for melanoma frequently and on a regular basis while using amantadine hydrochloride for any indication. Ideally, periodic skin examinations should be performed by appropriately qualified individuals (e.g., dermatologists). Other: The dose of amantadine hydrochloride may need careful adjustment in patients with congestive heart failure, peripheral edema, or orthostatic hypotension. Care should be exercised when administering amantadine hydrochloride to patients with a history of recurrent acute renal colic, or to patients with psychosis or severe psychiatric disorders not controlled by chemotherapeutic agents. Serious bacterial infections may begin with influenza-like symptoms or may cause death or occur as complications during the course of influenza. Amantadine hydrochloride has not been shown to prevent such complications. Information for Patients: Patients should be advised of the following information: Illnesses, eye pain, loss of vision, and/or impaired mental ability may occur. Gradually increase physical activity as the symptoms of Parkinson's disease improve. Avoid excessive alcohol usage, since it may increase the potential for CNS effects such as dizziness, confusion, lightheadedness and orthostatic hypotension. Avoid getting up suddenly from a sitting or lying position. It decreases or lightheadedness occurs, notify physician. Notify physician if mood, mental changes, swelling of extremities, difficulty urinating and/or shortness of breath occur. Do not take more medication than prescribed because of the risk of overdose. If there is no improvement in a few days, or if medication appears less effective after a few weeks, discuss with a physician. Consult physician before discontinuing medication. Seek medical attention immediately if it is suspected that an overdose of medication has been taken. Drug Interactions: Careful observation is required when amantadine hydrochloride is administered concurrently with central nervous system stimulants. Agents with anticholinergic properties may potentiate the anticholinergic-like side effects of amantadine hydrochloride. Combination of thioridazine has been reported to worsen the tremor in elderly patients with Parkinson's disease, however, it is not known if other phenothiazines produce a similar response. Combination of triamterene/hydrochlorothiazide resulted in a higher plasma amantadine hydrochloride concentration in a 64-year-old male taking amantadine hydrochloride 100 mg TID for Parkinson's disease.¹ It is not known which of the components of triamterene/hydrochlorothiazide contributed to the above effect or if related drugs produce a similar response. Combination of quinine or quinidine with amantadine hydrochloride was shown to reduce the renal clearance of amantadine hydrochloride by about 30%. The concurrent use of amantadine hydrochloride with low absorbed influenza vaccine (A/JA/Thailand/69) has not been evaluated. However, because of the potential for interference between these products, (A/JA) should not be administered within 2 weeks before or 48 hours after administration of amantadine hydrochloride, unless explicitly indicated. The concern about possible interference arises from the potential for antihist drugs to inhibit replication of the vaccine virus. Thioridazine, inactivated influenza vaccine can be administered at any time relative to use of amantadine hydrochloride. Cardiovascular: Widening/Deepening: Long-term (one year) animal studies designed to evaluate the cardioprotective potential of amantadine hydrochloride have not been performed. In several <i>in vitro</i> assays for gene mutations, amantadine hydrochloride did not increase the number of spontaneously induced mutations in four strains of <i>Salmonella typhimurium</i> (Ames Test) or in a mammalian cell line (Chinese Hamster Ovary cells) when incubations were performed either with or without a liver metabolic activation extract. Further, there was no evidence of chromosome damage observed in an <i>in vitro</i> test using freshly derived and stimulated human peripheral blood lymphocytes (with and without metabolic activation) or in an	<i>in vivo</i> mouse bone marrow micronucleus test (140 to 560 mg/kg; estimated human equivalent doses of 11.7 to 46.8 mg/kg based on body surface area comparison). Impairment of Fertility: The effect of amantadine hydrochloride on fertility has not been adequately tested, that is, in a study conducted under Good Laboratory Practice (GLP) and according to current recommended methodology. In a three litter, non-GLP, reproduction study in rats, amantadine hydrochloride at a dose of 32 mg/kg/day (equal to the maximum recommended human dose on a mg/m² basis) administered to both males and females slightly impaired fertility. There were no effects on fertility at a dose level of 10 mg/kg/day (or 0.3 times the maximum recommended human dose on a mg/m² basis); intermediate doses were not tested. Failed fertility has been reported during human <i>in vitro</i> fertilization (IVF) when the sperm donor ingested amantadine hydrochloride 2 weeks prior to, and during the IVF cycle. Pregnancy: Fetal/Neonatal Effects: The effect of amantadine hydrochloride on embryonic and post-natal development has not been adequately tested, that is, in studies conducted under Good Laboratory Practice (GLP) and according to current recommended methodology. However, in two non-GLP studies in rats in which females were dosed from 5 days prior to mating to Day 6 of gestation or on Days 7 to 14 of gestation, amantadine hydrochloride produced increases in embryonic death at an oral dose of 100 mg/kg for 3 times the maximum recommended human dose on a mg/m² basis. In the non-GLP rat study in which females were dosed on Days 7 to 14 of gestation, there was a marked increase in severe visceral and skeletal malformations at oral doses of 50 and 100 mg/kg (or 1.5 and 3 times, respectively, the maximum recommended human dose on a mg/m² basis). The no-effect dose for teratogenicity was 37 mg/kg (equal to the maximum recommended human dose on a mg/m² basis). The safety margins represented may not accurately reflect the risk considering the questionable quality of the study on which they are based. There are no adequate and well-controlled studies in pregnant women. Human data regarding teratogenicity after maternal use of amantadine hydrochloride is sparse. Fetal and placental malformations in normal karyotype fetuses in an infant exposed to amantadine hydrochloride during the first trimester of pregnancy (100 mg PO, for 7 days during the 1st and 7th week of gestation). Cardiovascular malformation (right ventricle with pulmonary stenosis) was associated with maternal exposure to amantadine hydrochloride (100 mg/kg) administered during the first 2 weeks of pregnancy. Amantadine hydrochloride should be used during pregnancy only if the potential benefit justifies the potential risk to the embryo or fetus. Breast Milk: Amantadine hydrochloride is secreted in human milk. Use is not recommended in nursing mothers. Pediatric Use: The safety and efficacy of amantadine hydrochloride in newborn infants and infants below the age of 1 year have not been established. Use in the Elderly: Because amantadine hydrochloride is primarily secreted in the urine, its accumulation in the plasma and in the body when renal function declines. Thus, the dose of amantadine hydrochloride should be reduced in patients with renal impairment and in individuals who are 65 years of age or older. The dose of amantadine hydrochloride may need reduction in patients with congestive heart failure, peripheral edema, or orthostatic hypotension (see DOSAGE AND ADMINISTRATION). ADVERSE REACTIONS The adverse reactions reported most frequently at the recommended dose of amantadine hydrochloride (5 to 10 mg) are: nausea, dizziness (lightheadedness), and insomnia. Less frequently (1 to 5%) reported adverse reactions are: depression, anxiety and irritability, hallucinations, confusion, anorexia, dry mouth, constipation, stools, libido reduction, peripheral edema, orthostatic hypotension, headache, somnolence, nervousness, dream abnormality, agitation, dry nose, diarrhea and fatigue. Infrequently (0.1 to 1%) occurring adverse reactions are: congestive heart failure, psychosis, urinary retention, dysphagia, skin rash, vomiting, weakness, slurred speech, euphoria, thinking abnormally, anorexia, hyperreflexia, hypertension, decreased libido, and visual disturbance, including parietal subarachnoid or other central opacity, corneal edema, decreased visual acuity, sensitivity to light, and optic nerve palsy. Rare (less than 0.1%) or occurring adverse reactions are: Instances of convulsion, leukopenia, neuropenia, acute/chronic dermatitis, ototoxicity symptoms, suicidal attempt, suicide, and suicidal ideation (see WARNINGS). Other adverse reactions reported during postmarketing experience with amantadine hydrochloride usage include: Neuroleptic Syndrome/Psychiatric: coma, stupor, delirium, hyperreflexia, hyperreflexia, aggression, behavior, paranoid ideation, muscle reaction, involuntary muscle contractions, gait abnormalities, parosmia, EEG changes, and tremor. Abrupt discontinuation may also precipitate delirium, agitation, delusions, hallucinations, paranoid reaction, sleep, anxiety, depression and disturbed speech. Cardiovascular: cardiac arrest, arrhythmias including malignant arrhythmias, hypertension, and tachycardia; Respiratory: acute respiratory failure, pulmonary edema, and tachypnea; Neuroleptic Syndrome: dysphagia; Hematologic: leukocytosis and agranulocytosis; Special Senses: keratitis, mydriasis, and corneal edema; Skin and Appendages: pruritus and dysphagia; Miscellaneous: neuroleptic malignant syndrome (see WARNINGS), allergic reactions including angioedema reactions, edema, and fever; Laboratory Test: elevated CPK, BUN, serum creatinine, alkaline phosphatase, LDK, bilirubin, SGOT, SGPT, and SALT.								
OVERDOSSAGE Cases have been reported from overdose with amantadine hydrochloride. The lowest reported acute lethal dose was 1 gram. Because some patients have attempted suicide by or overdosing with amantadine hydrochloride, precautions should be written for the smallest quantity available with good patient management. Acute toxicity may be attributable to the anticholinergic effects of amantadine hydrochloride. Drug overdoses has resulted in cardiac, respiratory, renal or central nervous system failure. Cardiac dysrhythmias include arrhythmias, bradycardia and hyperthermia. Pulmonary edema and respiratory distress (including adult respiratory distress syndrome - ARDS) have been reported; renal dysfunction including increased BUN, decreased creatinine clearance and renal insufficiency can occur. Central nervous system effects that have been reported include: insomnia, anxiety, agitation, aggressive behavior, hyperreflexia, hyperreflexia, stools, gut abnormality, tremor, confusion, disorientation, dysarthria, delirium, hallucinations, psychotic reactions, lethargy, somnolence and coma. Seizures have been reported in patients with prior history of seizure disorder. Hyperthermia has also been observed in cases where a drug overdose has occurred. There is no specific antidote for an overdose of amantadine hydrochloride. However, slowly administered intravenous physostigmine 1 and 2 mg doses in an adult at 1- to 2-hour intervals and 0.5 mg doses in a child at 5- to 10-minute intervals up to a maximum of 2 mg/hour have been reported to be effective in the control of central nervous system toxicity caused by amantadine hydrochloride. For acute overdosing, general supportive measures should be applied along with intravenous gastric lavage if induction of emesis. Fluids should be forced, and if necessary, given intravenously. The pH of the urine has been reported to influence the excretion rate of amantadine hydrochloride. Since the excretion rate of amantadine hydrochloride increases rapidly when the urine is acidic, the administration of urine acidifying drugs may increase the elimination of the drug from the body. The blood pressure, pulse, respiration and temperature should be monitored. The patient should be observed for pyrexia and convulsions. If required, sedation, and anticonvulsant therapy should be administered. The patient should be observed for the possible development of arrhythmias and hypotension. If required, appropriate antiarrhythmic and antihypertensive therapy should be given. ECG and cardiographic monitoring may be required after ingestion, since malignant tachyarrhythmias can appear after overdose. Care should be exercised when administering adverse agents, such as laxatives, to patients with an amantadine hydrochloride overdose, since the drug can synergize effects of amantadine hydrochloride has been reported to induce malignant arrhythmias. The blood electrolytes, urine pH and urinary output should be monitored. If there is no record of voiding, catheterization should be done. DOSAGE AND ADMINISTRATION The dose of amantadine hydrochloride may need reduction in patients with congestive heart failure, peripheral edema, orthostatic hypotension, or impaired renal function (see Dosage for Impaired Renal Function). Dosage for Prophylaxis and Treatment of Uncomplicated Influenza A Virus Illness: Adult: The adult daily dosage of amantadine hydrochloride is 200 mg (four teaspoons of oral solution) as a single daily dose. The daily dosage may be split into two teaspoons of oral solution twice a day. If central nervous system effects develop (insomnia - 4 day dosage, a split dosage schedule may reduce such complaints). In patients 65 years of age or older, the daily dosage of amantadine hydrochloride is 100 mg. A 100 mg daily dose has also been shown in experimental challenge studies to be effective in prophylaxis in healthy adults who are not at high risk for influenza related complications. However, it has not been demonstrated that a 100 mg daily dose is as effective as a 200 mg daily dose for prophylaxis, nor has the 100 mg daily dose been studied in the treatment of acute influenza illness. In recent clinical trials, the influence of central nervous system (CNS) side effects associated with the 100 mg daily dose was of or near the level of placebo. The 100 mg daily dose is recommended for persons who have a demonstrated intolerance to 200 mg of amantadine hydrochloride daily because of CNS or other liabilities. Pediatric Patients: 1 mg to 10 mg, of age. The total daily dose should be calculated on the basis of 2 to 4 mg/kg/day (4.4 to 8.8 mg/kg/day), but not to exceed 160 mg per day. 0 yrs. to 12 yrs. of age: The total daily dose is 200 mg given as two teaspoons of oral solution twice a day. The 100 mg daily dose has not been studied in this pediatric population. Therefore, there are no data which demonstrate that this dose is as effective as or is safer than the 200 mg daily dose in this patient population. Prophylaxis dosing should be initiated in anticipation of an influenza A virus outbreak and before or after contact with individuals with influenza A virus respiratory tract illness. Amantadine hydrochloride should be continued daily for at least 10 days following a known exposure. If amantadine hydrochloride is used chemoprophylactically in conjunction with inactivated influenza A virus vaccine until protective antibody responses develop, then it should be administered for 2 to 4 weeks after the vaccine has been given. When inactivated influenza A virus vaccine is unavailable or contraindicated, amantadine hydrochloride should be administered for the duration of known influenza. In the community because of repeated and unknown exposures. Treatment of influenza A virus illness should be started as soon as possible, preferably within 24 to 48 hours after onset of signs and symptoms, and should be continued for 24 to 48 hours after the disappearance of signs and symptoms.	Dosage for Parkinsonism: Adult: The usual dose of amantadine hydrochloride is 100 mg twice a day when used alone. Amantadine hydrochloride has an onset of activity usually within 48 hours. The initial dose of amantadine hydrochloride is 100 mg daily for patients with serious, associated medical illnesses or who are taking high doses of other antiparkinson drugs. After one to several weeks at 100 mg once daily, the dose may be increased to 100 mg twice daily, if necessary. Occasionally, patients whose responses are not optimal with amantadine hydrochloride at 200 mg daily may benefit from an increase up to 400 mg daily in divided doses. However, if such patients should be supervised closely by their physicians. Patients initially deriving benefit from amantadine hydrochloride not uncommonly experience a fall-off of effectiveness after a few months. Benefit may be regained by increasing the dose to 300 mg daily. Alternatively, temporary discontinuation of amantadine hydrochloride for several weeks, followed by reinitiation of the drug, may result in regaining benefit in some patients. A decision to use other antiparkinson drugs may be necessary. Dosage for Coccioidioid Tissue: Some patients who do not respond to anticholinergic antiparkinson drugs may respond to amantadine hydrochloride. When amantadine hydrochloride or anticholinergic antiparkinson drugs are each used with marginal benefit, concomitant use may produce additional benefit. When amantadine hydrochloride and levodopa are initiated concurrently, the patient can exhibit rapid therapeutic benefit. Amantadine hydrochloride should be held constant at 100 mg daily or twice daily while the daily dose of levodopa is gradually increased to optimal benefit. When amantadine hydrochloride is added to optimal well-tolerated doses of levodopa, additional benefit may result, including smoothing out the fluctuations in improvement which sometimes occur in patients on levodopa alone. Patients who require a reduction in their usual dose of levodopa because of development of side effects may possibly regain lost benefit with the addition of amantadine hydrochloride. Dosage for Drug-Induced Extrapyramidal Reactions: Adult: The usual dose of amantadine hydrochloride is 100 mg twice a day. Occasionally, patients whose responses are not optimal with amantadine hydrochloride at 200 mg daily may benefit from an increase up to 300 mg daily in divided doses. Dosage for Impaired Renal Function: Depending upon creatinine clearance, the following dosage adjustments are recommended: <table><tr><td>CREATININE CLEARANCE (mL/min/1.73m²)</td><td>AMANTADINE HYDROCHLORIDE DOSAGE</td></tr><tr><td>30 to 50</td><td>200 mg 1st day and 100 mg each day thereafter</td></tr><tr><td>15 to 30</td><td>200 mg 1st day followed by 100 mg on alternate days</td></tr><tr><td><15</td><td>200 mg every 7 days</td></tr></table> The recommended dosage for patients on hemodialysis is 200 mg every 7 days. HOW SUPPLIED Amantadine Hydrochloride Oral Solution, USP 50 mg/5 mL is a colorless to pale yellow, raspberry-flavored oral solution available in: 10 mL unit dose caps in bags of 10 (NDC 0116-4010-10) 1 Pint (512 mL) bottles (NDC 0116-4010-16) Store at 20° to 25° C (68° to 77° F), excursions permitted to 15° to 30° C (59° to 86° F). [See USP Controlled Room Temperature]. KEEP TIGHTLY CLOSED Dispense in a light container as defined in the USP, with a child-resistant closure (see USP). Do Not REFERENCES W.M. Wilson and A.H. Rajput, Amantadine-Dopamine Interaction, Can. Med. Assoc. J. 120:974-975, 1983. R.D. Cowley, Jr. Med. J. 286:516, 1978. R.D. Cowley, Jr. J. Pediatr. 85:144, 1974. Manufactured by: Xtrem Laboratories, Inc. 1230 E. Business Center Dr. Mount Pleasant, MI 48058 4010AMMHTS REV. 06-25	CREATININE CLEARANCE (mL/min/1.73m²)	AMANTADINE HYDROCHLORIDE DOSAGE	30 to 50	200 mg 1st day and 100 mg each day thereafter	15 to 30	200 mg 1st day followed by 100 mg on alternate days	<15	200 mg every 7 days
CREATININE CLEARANCE (mL/min/1.73m²)	AMANTADINE HYDROCHLORIDE DOSAGE								
30 to 50	200 mg 1st day and 100 mg each day thereafter								
15 to 30	200 mg 1st day followed by 100 mg on alternate days								
<15	200 mg every 7 days								

AMANTADINE

amantadine solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0116-4010
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
AMANTADINE HYDROCHLORIDE (UNII: M6Q1EO9TD0) (AMANTADINE - UNII:BF4C9Z1J53)		AMANTADINE HYDROCHLORIDE	50 mg in 5 mL

Inactive Ingredients			
Ingredient Name			Strength
METHYLPARABEN (UNII: A2I8C7HI9T)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)			

SORBITOL (UNII: 506T60A25R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0116-4010-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/05/2024	
2	NDC:0116-4010-41	100 in 1 CASE	11/05/2024	
2		10 in 1 TRAY		
2	NDC:0116-4010-10	10 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075060	11/05/2024	

Labeler - Xttrium Laboratories, Inc (007470579)

Registrant - Xttrium Laboratories, Inc (007470579)

Establishment

Name	Address	ID/FEI	Business Operations
Xttrium Laboratories, Inc.		007470579	manufacture(0116-4010)

Revised: 5/2025

Xttrium Laboratories, Inc